



Sphingosine 1 phosphate agonists (S1P); a potential agent to prevent acute lung injury in COVID-19

Sepideh Zununi Vahed¹, Shahram Ghiyasvand¹, Seyed Mahdi Hosseiniyan Khatibi¹, Bhargav Patel², Mohammadali Mohajel Shoja³, Ramin Tolouian⁴, Mohammadreza Ardalan¹

¹Kidney Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

²Department of Internal Medicine, University of Arizona, Tucson, AZ, USA

³Department of Surgery, University of Illinois at Chicago-Metropolitan Group Hospitals (UIC-MGH), Chicago, Illinois, USA

⁴Division of Nephrology, College of Medicine, University of Arizona, Tucson, AZ, USA

*Correspondence to

Prof. Mohammadreza Ardalan,
Email: ardalan34@yahoo.com

Received 20 May 2020

Accepted 14 June 2020

Published online 20 June 2020

Keywords: Cytokine storm, COVID-19, SARS-CoV-2, Fingolimod, FTY720-P, Acute respiratory distress syndrome

Abstract

SARS-CoV-2 is a worldwide pandemic, that has led to the morbidity and mortality of millions of people. This virus rapidly proliferates and destroys lung epithelial cells directly, which is worsened by a subsequent cytokine storm. This cytokine storm diffusely damages the alveolar barriers and leads to fibrin and fluid exudation, hyaline membrane formation, and infiltration of inflammatory cells into the lung causing acute respiratory distress syndrome (ARDS). To date, there exists no medication to treat SARS-CoV-2 infection and novel new therapeutics are still being explored to prevent or limit the damage to the lung. Sphingosine 1-phosphate (S1P) is an effective bioactive lipid mediator and its related signaling pathways are vital for endothelial cell integrity. Stabilizing the pulmonary endothelial barrier and decreasing the inflammatory infiltrate by S1P analogs such as Fingolimod (FTY720-P) would be a new therapeutic approach for the hindrance of pulmonary exudation and subsequent ARDS.

Introduction

Severe acute respiratory syndrome coronavirus virus 2 (SARS-CoV-2) targets the respiratory epithelial cells, which leads to a severe respiratory presentation in 15% to 20% of patients. This virus rapidly proliferates, destroys the epithelial cells, and triggers the immune system causing a cytokine storm. This cytokine storm diffusely damages the alveolar barriers and leads to fibrin and fluid exudation, hyaline membrane formation, and infiltration of inflammatory cells into the lung causing acute respiratory distress syndrome (ARDS). Morbidity and mortality of SARS-CoV-2 infection are directly related to this dysregulated immune response (1). Researchers identified Sphingosine 1-phosphate (S1P), a membrane sphingolipid metabolite, as a biologically active lipid-signaling mediator in the 1990s (2). S1P and its receptors have a pivotal role in the preservation of vascular endothelial cell barrier integrity and the regulation of immune cells' diapedesis. S1P binds to a five-membrane related G-protein coupled receptors (GPCRs) including S1PR 1, -2, and -3 which are present in the cardiovascular system, while

Key point

Stabilizing the endothelial barrier and decreasing the inflammatory infiltrate by a sphingosine 1-phosphate analog (Fingolimod, FTY720) would be a novel therapeutic strategy towards hindering pulmonary exudation.

the S1P1R is the dominant receptor in the lung. S1P binds to S1P1R on the endothelial cell to improve the integrity of endothelial barrier and restrict the effector lymphocytes migration toward the inflamed area and infiltration to the alveolar space (3). The activation of S1P1R additionally displays a cytokine blunting activity and attenuates IL-6 secretion (4). In the following paragraphs, we will concisely discuss the existing science on S1P signaling in the lung.

Pathology of the SARS-CoV-2

SARS-CoV-2 was a novel and sudden disease outbreak, therefore there are very few lung histopathology reports. In the lung histopathology studies of two asymptomatic patients, who underwent pulmonary surgery because of another reason, but were later

Citation: Zununi Vahed S, Ghiyasvand S, Hosseiniyan Khatibi SM, Patel B, Mohajel Shoja M, Tolouian R, et al. Sphingosine 1 phosphate agonists (S1P); a potential agent to prevent acute lung injury in COVID-19. *Immunopathol Persa.* 2021;7(1):e03. DOI:10.34172/ipp.2021.03.



seropositive, revealed pneumocyte hyperplasia with patchy inflammatory cellular infiltration without any hyaline membrane formation. We can consider these findings as early-stage involvements (5). Postmortem autopsy findings in a 50-year-old patient who died due to COVID-19 respiratory failure resembled those seen in Middle Eastern respiratory syndrome (MERS) and SARS. The important findings were prominent lymphocytic interstitial infiltration dominated by cytotoxic CD8 and Th17 cells. These findings show the importance of inflammatory cells in lethal cases of SARS-CoV-2 infection (6).

Role of S1P in endothelial barrier regulation

Over the past decade, several studies have validated that S1PR1 signaling is imperative for the enhancement of lung endothelial barrier function (7-9). Enhancing the effects of S1P on endothelial barrier are ascribed after the activation of S1PR1, which subsequently activates downstream signaling through Rho GTPases and then stabilizes the endothelial actin cytoskeleton (10). S1P mediated localization of beta-catenin and VE-cadherin at the sites of endothelial cells' adherent junctions maintains vascular integrity (11).

The intracellular myosin light chain phosphorylation and the rearrangement of adherent junction proteins in the lung endothelium cells have been proposed as possible S1P-induced barrier function improvement (12-15). In an *in vivo* model of ventilator-associated lung damage, S1P analog stabilizes endothelial cytoskeleton, cell-matrix adherence and tightens the inter-cellular junction to prevent alveolar exudation (16-18).

Role of S1P activation in viral-induced lung injuries

H1N1 viral infection causes a cytokine storm, whose major target is endothelial cells. Based on experimental findings in a mouse model of H1N1 infection, intensified endothelial S1PR1 signaling blunts the cytokine-induced inflammatory cell infiltration (19). A S1PR1 agonist preserves the lung tissue by diminishing the endothelial cells response to a high cytokine state. In H1N1 influenza murine models (2009), S1PR1 agonists showed a decrease of more than 80% of deaths compared to 50% protection by oseltamivir, an antiviral neuraminidase inhibitor. The combination therapy of these drugs had optimal protection up to 96% (20). The immune-modulatory strategies using S1P analog had a promising outcome in influenza infection (21). In another study, the combined usage of S1PR1 agonist; oseltamivir and CYM-5442 had the greatest protection against the influenza-induced lung injury (22).

FTY720

FTY720 (Fingolimod) is a sphingosine analog and agonist of S1P receptors. Sphingosine kinase (SphK) phosphorylates FTY720 to its active form FTY720-P.

In 2010, the Food and Drug Administration of United States (FDA), approved FTY720 as a treatment of multiple sclerosis (23). Several murine and canine studies have showed meaningful reduction in pulmonary damage after treatment with FTY720 or Sph 1-P (16,24). Additionally, another study showed FTY720 inhibits lymphocyte-mediated airway inflammation (25) and ameliorates central nervous system-induced inflammatory cell recruitment in experimental autoimmune encephalomyelitis (26). Protective impacts of S1P analogous and FTY720 has been described in vascular endothelial growth factor (VEGF)-mediated models of vascular leakage (27,28). Recent findings also support the role of FTY720 on the endothelial barrier enhancement and subsequent suppression of trans-endothelial migration of inflammatory cells (29-31).

Proposed hypothesis

The clinical accessibility of S1P agonist, FTY720, makes it a striking therapeutic option. The 0.5 mg daily oral administration of FTY720 for three consecutive days within the first 72 hours of acute ischemic stroke is associated with a noteworthy reduction of ischemic lesion expansion and its administration protects the brain from the intense lymphocytes infiltration that happens after stroke-induced cytokine storm (32); we can make an analogy between those findings and maybe the same course of FTY720 (Fingolimod protocol) could be a useful scheme in high-risk individuals with COVID-19 infection. The early start of S1P agonist may prevent the pulmonary infiltration and functional deterioration.

Conclusion

S1P is an effective bioactive lipid mediator and its related signaling pathways might be a crucial area for new therapeutics for SARS-CoV-2. Stabilizing pulmonary endothelial barrier and decreasing the inflammatory infiltrate by S1P analog/FTY720 is a novel therapeutic approach towards hindering pulmonary exudation, a major cause of morbidity and mortality in SARS-CoV-2 patients (Figure 1).

Authors' contribution

SZV and MRA developed the idea and revised the manuscript. ShG and SMHK contributed in preparing original draft. MMS and RT revised the manuscript. BP revised and edited the manuscript.

Conflicts of interest

No conflict of interest.

Ethical considerations

Ethical issues (including data fabrication, double publication, and plagiarism) have been detected by the authors.

Funding/Support

None.

References

- Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical




Figure 1. Sphingosine 1 phosphate agonist is a promising plan against SARS-CoV-2-induced pulmonary damage.

ACE2 is identified as the functional receptor for SARS-CoV-2. ACE2 is highly present on human alveolar epithelial cells (33), its high expression is identified in type II alveolar cells facilitating the replication of coronavirus in the lung (34, 35). The injured cells promote an innate inflammation in the lungs mediated by pro-inflammatory macrophages (6). S1P is an effective bioactive lipid mediator and its related signaling pathways are crucial for endothelial integrity. As a S1P analogs, FTY720-P (Fingolimod) binding to S1P1R on the endothelial cell enhances the endothelial barrier integrity and limits the migration of effector lymphocytes toward the inflamed area. It can also decrease the fluid extravasation and cytokine storm in the alveolar space (3). ACE2: angiotensin-converting enzyme 2, S1P: Sphingosine 1-phosphate, Sph: sphingosine, MQ: macrophage,

features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497-506. doi: 10.1016/s0140-6736(20)30183-5

- Ghosh TK, Bian J, Gill DL. Intracellular calcium release mediated by sphingosine derivatives generated in cells. *Science*. 1990;248:1653-6. doi: 10.1126/science.2163543.
- Rosen H, Goetzl EJ. Sphingosine 1-phosphate and its receptors: an autocrine and paracrine network. *Nat Rev Immunol*. 2005;5: 560-70. doi:10.1038/nri1650.
- Zhao Y, Gorshkova IA, Berdyshev E, He D, Fu P, Ma W, et al. Protection of LPS-induced murine acute lung injury by sphingosine-1-phosphate lyase suppression. *Am J Respir Cell Mol Biol*. 2011;45:426-35. doi: 10.1165/rcmb.2010-0422.OC.
- Tian S, Hu W, Niu L, Liu H, Xu H, Xiao SY. Pulmonary Pathology of Early-Phase 2019 Novel Coronavirus (COVID-19) Pneumonia in Two Patients With Lung Cancer. *J Thorac Oncol*. 2020. doi: 10.1016/j.jtho.2020.02.010.
- Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med*. 2020. doi: 10.1016/s2213-2600(20)30076-x.
- Peng X, Hassoun PM, Sammani S, McVerry BJ, Burne MJ, Rabb H, et al. Protective effects of sphingosine 1-phosphate in murine endotoxin-induced inflammatory lung injury. *Am J*

Respir Crit Care Med. 2004;169(11):1245-51. doi: 10.1164/rccm.200309-1258OC.

- Wang L, Dudek SM. Regulation of vascular permeability by sphingosine 1-phosphate. *Microvasc Res*. 2009;77:39-45. doi: 10.1016/j.mvr.2008.09.005.
- Sammani S, Moreno-Vinasco L, Mirzapioazova T, Singleton PA, Chiang ET, Evenoski CL, et al. Differential effects of sphingosine 1-phosphate receptors on airway and vascular barrier function in the murine lung. *Am J Respir Cell Mol Biol*. 2010;43:394-402. doi: 10.1165/rcmb.2009-0223OC
- Garcia JG, Liu F, Verin AD, Birukova A, Dechert MA, Gerthoffer WT, et al. Sphingosine 1-phosphate promotes endothelial cell barrier integrity by Edg-dependent cytoskeletal rearrangement. *J Clin Invest*. 2001;108:689-701. doi: 10.1172/jci12450.
- Lee MJ, Thangada S, Claffey KP, Ancellin L, Liu CH, Kluk M, et al. Vascular endothelial cell adherens junction assembly and morphogenesis induced by sphingosine-1-phosphate. *Cell*. 1999;99:301-12. doi: 10.1016/s0092-8674(00)81661-x.
- Dudek SM, Jacobson JR, Chiang ET, Birukov KG, Wang P, Zhan X, et al. Pulmonary endothelial cell barrier enhancement by sphingosine 1-phosphate: roles for cortactin and myosin light chain kinase. *J Biol Chem*. 2004;279:24692-700. doi:10.1074/jbc.M313969200.
- Sun X, Shikata Y, Wang L, Ohmori K, Watanabe N, Wada J, et al. Enhanced interaction between focal adhesion and adherens junction proteins: involvement in sphingosine 1-phosphate-induced endothelial barrier enhancement. *Microvasc Res*. 2009;77:304-13. doi: 10.1016/j.mvr.2008.12.004
- Abbasi T, Garcia JG. Sphingolipids in lung endothelial biology and regulation of vascular integrity. *Handb Exp Pharmacol*. 2013;(216):201-26. doi: 10.1007/978-3-7091-1511-4_10.
- Xiong Y, Hla T. S1P control of endothelial integrity. *Curr Top Microbiol Immunol*. 2014;378:85-105. doi: 10.1007/978-3-319-05879-5_4
- McVerry BJ, Peng X, Hassoun PM, Sammani S, Simon BA, Garcia JG. Sphingosine 1-phosphate reduces vascular leak in murine and canine models of acute lung injury. *Am J Respir Crit Care Med*. 2004;170:987-93. doi: 10.1164/rccm.200405-684OC.
- Teijaro JR, Walsh KB, Cahalan S, Fremgen DM, Roberts E, Scott F, et al. Endothelial cells are central orchestrators of cytokine amplification during influenza virus infection. *Cell*. 2011;146:980-91. doi:10.1016/j.cell.2011.08.015.
- Walsh KB, Teijaro JR, Rosen H, Oldstone MB. Quelling the storm: utilization of sphingosine-1-phosphate receptor signaling to ameliorate influenza virus-induced cytokine storm. *Immunol Res*. 2011;51:15-25. doi: 10.1007/s12026-011-8240-z.
- Teijaro JR, Walsh KB, Long JP, Tordoff KP, Stark GV, Eisfeldt AJ, et al. Protection of ferrets from pulmonary injury due to H1N1 2009 influenza virus infection: immunopathology tractable by sphingosine-1-phosphate 1 receptor agonist therapy. *Virology*. 2014;452-453:152-7. doi: 10.1016/j.virol.2014.01.003.
- Oldstone MB, Teijaro JR, Walsh KB, Rosen H. Dissecting influenza virus pathogenesis uncovers a novel chemical approach to combat the infection. *Virology*. 2013;435:92-101. doi: 10.1016/j.virol.2012.09.039
- Walsh KB, Teijaro JR, Wilker PR, Jatzek A, Fremgen DM, Das SC, et al. Suppression of cytokine storm with a sphingosine analog provides protection against pathogenic influenza virus. *Proc Natl Acad Sci U S A*. 2011;108:12018-23. doi: 10.1073/pnas.1107024108.
- Zhao J, Zhu M, Jiang H, Shen S, Su X, Shi Y. Combination of sphingosine-1-phosphate receptor 1 (S1PR1) agonist and antiviral drug: a potential therapy against pathogenic influenza virus. *Sci Rep*. 2019;9:5272. doi: 10.1038/s41598-019-

- 41760-7.
23. Pitman M, Woodcock J, Lopez A, Pitson S. Molecular targets of FTY720 (fingolimod). *Curr Mol Med*. 2012;12:1207-9. doi: 10.2174/156652412803833599
 24. Liu HB, Cui NQ, Wang Q, Li DH, Xue XP. Sphingosine-1-phosphate and its analogue FTY720 diminish acute pulmonary injury in rats with acute necrotizing pancreatitis. *Pancreas*. 2008;36(3):e10-5. doi: 10.1097/MPA.0b013e31815f3905
 25. Sawicka E, Zuany-Amorim C, Manlius C, Trifilieff A, Brinkmann V, Kemeny DM, et al. Inhibition of Th1- and Th2-mediated airway inflammation by the sphingosine 1-phosphate receptor agonist FTY720. *J Immunol*. 2003;171:6206-14. doi: 10.4049/jimmunol.171.11.6206
 26. Fujino M, Funeshima N, Kitazawa Y, Kimura H, Amemiya H, Suzuki S, et al. Amelioration of experimental autoimmune encephalomyelitis in Lewis rats by FTY720 treatment. *J Pharmacol Exp Ther*. 2003;305:70-7. doi: 10.1124/jpet.102.045658
 27. Sanchez T, Estrada-Hernandez T, Paik JH, Wu MT, Venkataraman K, Brinkmann V, et al. Phosphorylation and action of the immunomodulator FTY720 inhibits vascular endothelial cell growth factor-induced vascular permeability. *J Biol Chem*. 2003;278:47281-90. doi: 10.1074/jbc.M306896200
 28. English D, Kovala AT, Welch Z, Harvey KA, Siddiqui RA, Brindley DN, et al. Induction of endothelial cell chemotaxis by sphingosine 1-phosphate and stabilization of endothelial monolayer barrier function by lysophosphatidic acid, potential mediators of hematopoietic angiogenesis. *J Hematother Stem Cell Res*. 1999;8:627-34. doi: 10.1089/152581699319795
 29. Imeri F, Fallegger D, Zivkovic A, Schwalm S, Enzmann G, Blankenbach K, et al. Novel oxazolo-oxazole derivatives of FTY720 reduce endothelial cell permeability, immune cell chemotaxis and symptoms of experimental autoimmune encephalomyelitis in mice. *Neuropharmacology*. 2014;85:314-27. doi: 10.1016/j.neuropharm.2014.05.012
 30. Camp SM, Chiang ET, Sun C, Usatyuk PV, Bittman R, Natarajan V, et al. Pulmonary Endothelial Cell Barrier Enhancement by Novel FTY720 Analogs: Methoxy-FTY720, Fluoro-FTY720, and beta-Glucuronide-FTY720. *Chem Phys Lipids*. 2016;194:85-93. doi: 10.1016/j.chemphyslip.2015.10.004
 31. Hemdan NY, Weigel C, Reimann CM, Graler MH. Modulating sphingosine 1-phosphate signaling with DOP or FTY720 alleviates vascular and immune defects in mouse sepsis. *Eur J Immunol*. 2016;46:2767-2777. doi: 10.1002/eji.201646417.
 32. Fu Y, Zhang N, Ren L, Yan Y, Sun N, Li YJ, et al. Impact of an immune modulator fingolimod on acute ischemic stroke. *Proc Natl Acad Sci U S A*. 2014;111:18315-20. doi: 10.1073/pnas.1416166111.
 33. Hamming I, Timens W, Bulthuis ML, Lely AT, Navis G, van Goor H. Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis. *J Pathol*. 2004;203: 631-7. doi: 10.1002/path.1570.
 34. Zou X, Chen K, Zou J, Han P, Hao J, Han Z. Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Front Med*. 2020. doi: 10.1007/s11684-020-0754-0.
 35. Zhang H, Penninger JM, Li Y, Zhong N, Slutsky AS. Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive Care Med*. 2020. doi: 10.1007/s00134-020-05985-9.