



# Potential factors related to post-COVID-19 persistent symptoms; Khorshid COVID cohort study

Ramin Sami<sup>1</sup>, Zahra Khani Khoozani<sup>2</sup>, Marjan Mansourian<sup>3</sup>, Zohre Naderi<sup>2</sup>, Mehdi Karami<sup>4</sup>, Mina Nickpour<sup>1</sup>, Arash Toghiani<sup>2</sup>, Ali Gharavinia<sup>5</sup>, Nastaran Sadat Hosseini<sup>2</sup>, Mehrnegar Dehghan<sup>2</sup>, Midia Babahajiani<sup>6</sup>, Nafiseh Shokri-Mashhadi<sup>7</sup>

<sup>1</sup>Department of Internal Medicine, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>2</sup>Department of Internal Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>3</sup>Pediatric Cardiovascular Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>4</sup>Department of Radiology, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>5</sup>Department of Family and Emergency Medicine, School of Medicine, Azad University of Medical Science, Isfahan, Iran.

<sup>6</sup>Student Research Committee, Vice Chancellor for Research and Technology, Kurdistan University of Medical Sciences, Sanandaj, Iran

<sup>7</sup>Department of Clinical Nutrition, School of Nutrition and Food Science, Food Security Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

## \*Correspondence to

Nafiseh Shokri-Mashhadi,

Email: n.shokri@nutr.mui.ac.ir,

nafiseh.shokri@yahoo.com

Received 1 Jan. 2021

Accepted 9 Feb. 2021

Published online 17 Feb. 2021

**Keywords:** COVID-19, Post-discharge, Follow up, Persistent symptoms

## Abstract

**Introduction:** Recently, a high percentage of post-discharge patients have reported persistent symptoms after COVID-19 recovery. However, there is limited comprehensive information regarding nature's symptoms and association with potential factors.

**Objectives:** We aimed to assess the inpatients' characteristics and the prevalence of COVID-19 symptoms at admission and four weeks after recovery to recognize the potential factors related to the persistent symptoms.

**Patients and Methods:** We assessed the medical information of 262 severe and 546 non-severe COVID-19 inpatients at admission, and during the first and four weeks after post-discharge from the Khorshid COVID Cohort (KCC) study.

**Results:** Cough, dyspnea, and fatigue were the most reported symptoms at admission, and continuously were declined over the time (all  $P < 0.01$ ). However, the complaint of weight loss was increased during follow-up ( $P = 0.01$ ). Older age ( $P < 0.0001$ ), male gender ( $P = 0.02$ ), administration of hydroxychloroquine ( $P = 0.017$ ), and the interval time from illness onset to visit hospital ( $P < 0.0001$ ) augmented the remaining respiratory symptoms risk. Additionally, more length of hospital stay correlated to the lower risk of persistent constitutional symptoms ( $P < 0.05$ ).

**Conclusion:** This study points out the greater rehabilitation needs and management of persistent symptoms, in particular cough, dyspnea, fatigue and weight loss, and their related factors.

**Citation:** Sami R, Khani khoozani Z, Mansourian M, Naderi Z, Karami M, Nickpour M, Toghiani A, Gharavinia A, Hosseini NS, Dehghan M, Babahajiani M, Shokri-Mashhadi N. Potential factors related to post-COVID-19 persistent symptoms; Khorshid-COVID cohort study. Immunopathol Persa. 2023;9(2):e22223x. DOI:10.34172/ipp.2022.22223.

## Introduction

Recently, with the increase in the number of hospitalization of severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) cases (1), new problems appeared in the light of following up of patients after recovery (2,3). The high proportion of COVID-19 survivors discharged from the hospital might experience persistent constitutional and respiratory post-discharge symptoms and abnormal radiological outcomes (2,4). Fatigue, dyspnea and psychotic symptoms were the most prevalent symptoms after discharge from the hospital (5, 6).

Recognize the nature of remaining symptoms after recovery helps healthcare services to plan the rehabilitation needs in survivors of COVID-19 infection. However, there is limited comprehensive information

## Key point

This study highlighted the emergency need for management of persistent symptoms, in particular, cough, dyspnea, fatigue, and weight loss and their modifiable related factors in order to control post-discharge COVID-19 infection.

regarding the prevalence and factors contributing to the remaining symptoms after discharge from the hospital, and their association with the patient's characteristics (7).

## Objectives

We aimed to assess the patients' characteristics and the prevalence of COVID-19 symptoms during care at the hospital and four weeks after recovery to recognize the potential



factors related to the persistent symptoms.

## Patients and Methods

### Data extraction

The following information of confirmed COVID-19 patients ( $\geq 18$  years old) was extracted from the Khorshid COVID Cohort (KCC) study, from February 2020 to April 2020 (3). All adult patients who had coronavirus symptoms with typical CT scan lesions or positive real-time reverse transcriptase-polymerase chain reaction (RT-PCR) responses were selected by the census method and enrolled in this study. We excluded recorded data about death cases and patients who refused to participate. The disease was defined as severe if the respiratory frequency was more than 30 breaths per minute or blood oxygen saturation was less than 88%. We considered other cases as non-severe COVID-19 patients. Besides collecting some general information about demographics, pre-existing comorbidities, date of onset of symptoms, and COVID-19 diagnosis, additional follow-up information concerning the presence of post-discharge persistent symptoms was collected at the first and fourth week by two trained physicians using a short phone questionnaire.

### Statistical analysis

We performed the  $\chi^2$  test for categorical data and Fisher's exact test for continuous variables to compare the variables regarding illness severity (severe, non-severe). The categorical data differences between paired samples were estimated using McNamara's test. A value of  $P < 0.05$  was considered statistically significant. Association between the persistence symptoms and other characteristics of patients was estimated by generalized estimating equation analysis through the four-week follow-up. Two repeated measurements of each patient (the first and fourth weeks after discharge) were accounted for an exchangeable correlation structure and robust variance estimation.

## Results

Of 1350 inpatients, characteristics of 876 recovered COVID-19 patients were evaluated, of whom the information of 10 patients who died and 58 cases who denied to follow-up was excluded from the study. In total, 808 patients completed the first week of follow-up and 806 patients remained to the fourth week. The median age in non-severe and severe patients were 55 years (44-66), and 63 years (53-73), respectively ( $P < 0.001$ ). Around 58.9% of all patients were male (Table 1). The time from illness onset to visit hospital ranged from 4-10 days, and the median range of CT score was 10 (6-13). The ratio of severe cases of COVID-19 under active quarantine was more than non-severe patients ( $P = 0.09$ ). The median levels of C-reactive protein (CRP) at admission were 29 (13-53) mg/L. Most patients had at least two pre-existing co-morbidities previously (median [IQR], 2 (1-4).

### Symptoms at admission

There was no significant difference in respiratory symptoms between severe and non-severe subgroups of COVID-19 infection (Figure 1A, B). The most prevalent impaired outcomes were cough (76%), fever (70.6), dyspnea (65%), and fatigue (57%).

### Symptoms at follow-up

In the first week of follow-up, the most reported symptoms were chest pain (78.6%), cough (25.2%), and dyspnea (20.6%). Inversely, cough ( $P < 0.001$ ), chest pain ( $P < 0.006$ ), and anosmia ( $P < 0.045$ ) were significantly more prevalent in non-severe patients than those in severe ones. Besides, all patients had more complaints of chest pain in the first week compared with admission, 48.6% in the first week versus 35% at admission. However, the sore throat symptom disappeared entirely in all cases (Figure 1A, B).

After four weeks, the presence of dyspnea persisted in approximately 15% of non-severe cases and it increased in patients who experienced severe COVID-19 disease over time (2%). Moreover, most post-discharge patients had complaints of chest pain in the first (41% in severe and 52% in non-severe,  $P < 0.006$ ) and fourth weeks follow-up (20% in severe and 12% in non-sever patients,  $P < 0.001$ ). After four weeks, there were non-significant differences in severe and non-severe COVID-19 patients regarding dyspnea, 20.6% and 14.86%, respectively ( $P = 0.056$ ). There were no significant variances concerning fatigue in severe and non-severe post-discharge individuals throughout the first week, 10.2% and 12.2%, respectively ( $P = 0.429$ ). The gastrointestinal disorder symptoms had low-prevalence during the follow-up. Inversely, anosmia appeared in the fourth week of follow-up in severe (4.4%) and non-sever patients (5%).

### Prediction of COVID-19 persistent symptoms risk

Besides, older age (odds ratio [OR]: 1.02; 95% CI: 1.019-1.045,  $P < 0.0001$ ), male gender (OR: 1.32; 95% CI: 1.034-1.705,  $P = 0.02$ ), administration of hydroxychloroquine (OR, 1.35; 95% CI, 1.056-1.738,  $P = 0.017$ ), and more interval time to visit hospital (OR: 2.31; 95% CI, 2.02-2.55,  $P < 0.0001$ ) increased the risk of the persistent respiratory symptoms. However, there was no association between the results of CT score or disease severity and the risk of remaining symptoms. The length of hospital stay and hydroxychloroquine administration significantly decreased the risk of remaining constitutional symptoms (OR: 0.96; 95% CI: 9.37-9.87 and OR: 0.74; 95% CI: 0.56-0.99, all  $P < 0.05$ , respectively). Interestingly, laboratory tests, including CRP, neutrophil-to-lymphocyte ratio (NLR), lactate dehydrogenase (LDH), levels, did not relate to the risk of persistent symptoms.

## Discussion

Our findings indicated that as per previous study, cough, dyspnea and chest pain are the most frequent respiratory

**Table 1.** Baseline demographic and clinical characteristics of COVID-19 inpatients on admission and 1st and 4th weeks after discharge

	All cases (N = 808)	Severity		P value
		Non-Severe 546 (67.6%)	Severe 262 (32.4%)	
Age, Median (IQR) years	58 (46-69)	55 (44-66)	63 (53-73)	<0.001
Age groups (y), No. (%)				
20-44	175 (21.7%)	145 (26.5%)	30 (11.45%)	
45-64	372 (46%)	254 (46.5%)	118 (45.03%)	
≥65	261 (32.3%)	147 (26.92%)	114 (43.51%)	
Male, No. (%)	476 (58.9%)	320 (58.60%)	156 (59.54%)	0.801
Travel to high risk place	128 (15.8%)	92 (16.8%)	36 (13.74%)	0.253
Quarantine n (%)	685 (84.8%)	455 (83.3%)	230 (87.78%)	0.09
Days from illness onset to visit hospital	7 (4-10)	7 (4-10)	7 (4-10)	0.550
Upper respiratory symptom				
On admit	263 (32.5%)	363 (66.4%)	193 (73.66%)	
1 <sup>st</sup> week*	10 (1.2%)	6 (1.1%)	4 (1.52%)	0.737
4 <sup>th</sup> week**	605 (74.8%)	422 (77.28%)	183 (66.84%)	0.005
Lower respiratory symptom				
On admit	724 (89.6%)	494 (90.47%)	230 (87.78%)	0.044
1 <sup>st</sup> week	570 (70.54%)	408 (74.72%)	162 (61.83%)	<0.001
4 <sup>th</sup> week	260 (32.17%)	160 (29.30%)	100 (38.16%)	0.023
Constitutional symptom				
On admit	766 (94.8%)	516 (94.50%)	250 (95.41%)	0.445
1 <sup>st</sup> week	176 (21.7%)	114 (20.87%)	62 (23.66%)	0.472
4 <sup>th</sup> week	195 (24.13%)	130 (23.80%)	65 (24.80%)	0.901
Gastrointestinal symptom				
On admit	453 (56.06%)	309 (56.59%)	144 (54.96%)	0.425
1 <sup>st</sup> week	49 (6.0%)	34 (6.22%)	15 (5.72%)	0.718
4 <sup>th</sup> week	25 (3.09%)	17 (3.1%)	8 (3.05%)	0.917
Neurologic symptom				
On admit	364 (45.04%)	273 (50.0%)	91 (34.73%)	<0.001
1 <sup>st</sup> week	29 (3.58%)	25 (4.57%)	4 (1.52%)	0.025
4 <sup>th</sup> week	40 (4.95%)	26 (4.76%)	14 (5.34%)	0.779
Mortality at 4 <sup>th</sup> week	16 (1.98%)	3 (.54)	13 (4.96%)	<0.001

\*1<sup>st</sup> week after follow up; \*\*4<sup>th</sup> week after follow up.


prolonged symptoms of COVID-19 infection (2). Although the post-discharge chest pain can be associated with several conditions (including tracheitis, pleuritis, pericarditis, GERD (gastroesophageal reflux disease), and musculoskeletal disorders) (8, 9), in some cases, there is no particular reason describing this condition.

Concerning post-discharge fatigue, some studies recently indicated that chronic fatigue syndrome could be as a long-term adverse effect of COVID-19 and caused by neurological exhaustion (10,11). Regarding anosmia manifestation after four weeks, we hypothesized that it could be a delayed symptom of COVID-19 infection, recognized as a hallmark of COVID-19 (12). This symptom may indicate the neuroinvasion of COVID-19 that could result in chronic neurodegenerative disease along with

hospitalization in high-risk patients due to the exacerbated cytokines release and neuroinflammation (13). The relationship between prolonged respiratory symptoms and protease inhibitors-based regimens indicated before (14). It seems that patients with more length of hospital stay had lower chances of having persistent constitutional post-discharge symptoms. Nevertheless, owing to coronavirus conflicting outcomes and limited data about post-discharge symptoms, we could not precisely realize the results underlying this condition.

### Conclusion

This observational study points out the emergency need to cure and manage persistent symptoms, in particular cough, dyspnea, fatigue, and weight loss and their



**Figure 1.** (A) Clinical characteristics of non-severe patients at admission, first and fourth weeks after discharge. B) Clinical characteristics of severe patients at admission, first and fourth weeks after discharge.

modifiable related factors, including hydroxychloroquine-based treatments, and the interval time from illness onset to visit the hospital, in order to control post-discharge COVID-19 infection in all patients.

### Limitations of the study

Our study has some limitations. Our survey does not include post-discharge pulmonary function tests. Moreover, our analysis relied on the follow-up information of self-reported questionnaires, which may subject to self-reporting and responding bias. We also specifically studied the experience of hospitalized people. Therefore, the community-based study should be designed to assess post-discharge symptoms of Covid-19 patients in the community.

### Authors' Contribution

**Conceptualization:** RS, NSM

**Methodology:** RS, ZN, MK ZKK, MN, AG, AT, NSH, MM and NS

**Validation:** RS, NSM,

**Formal analysis:** RS, NSM

**Investigation:** RS,

**Resources:** RS, MD, MB and NS

**Data curation:** RS, MD, MB and NS

**Writing—original draft:** RS, NSM

**Writing—review and editing:** RS, NSM

**Visualization:** MD, MB

**Supervision:** RS, ZN, MK ZKK, MN, AG, AT, NSH, MM and NS.

**Project administration:** RS, NSM.

### Conflicts of interest

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Ethical issues

The research followed the tenets of the Declaration of Helsinki. The Ethics Committee of Isfahan University of Medical Sciences approved this study (IR.MUI.MED.REC.1399.680). Accordingly, written informed consent was obtained from all the respondents at the start of the study. This study was extracted from the internal medicine residency thesis of Zahra Khani Khoozani at Isfahan University of Medical Sciences (Proposal #399637). Besides, ethical issues (including plagiarism, data fabrication, double publication) have been completely observed by the authors.

### Funding/Support

This study was supported by Isfahan University of medical science (Grant# 399637).

## References

1. Asyary A, Veruswati M. Sunlight exposure increased Covid-19 recovery rates: A study in the central pandemic area of Indonesia. *Sci Total Environ.* 2020 Aug 10;729:139016. doi: 10.1016/j.scitotenv.2020.139016.
2. Carfi A, Bernabei R, Landi F; Gemelli Against COVID-19 Post-Acute Care Study Group. Persistent Symptoms in Patients After Acute COVID-19. *JAMA.* 2020;324(6):603-605. doi: 10.1001/jama.2020.12603.
3. Sami R, Soltaninejad F, Amra B, Naderi Z, Haghjooy Javanmard S, et al. A one-year hospital-based prospective COVID-19 open-cohort in the Eastern Mediterranean region: The Khorshid COVID Cohort (KCC) study. *Plos One.* 2020;15:e0241537. doi: 10.1371/journal.pone.0241537.
4. Kazama I. Stabilizing mast cells by commonly used drugs: a novel therapeutic target to relieve post-COVID syndrome? *Drug Discov Ther.* 2020;14(5):259-261. doi: 10.5582/ddt.2020.03095.
5. Goertz YMJ, Van Herck M, Delbressine JM, Vaes AW, Meys R, Machado FVC, et al. Persistent symptoms 3 months after a SARS-CoV-2 infection: the post-COVID-19 syndrome? *ERJ Open Res.* 2020;6:00542-2020. doi: 10.1183/23120541.00542-2020.
6. Lim ST, Janaway B, Costello H, Trip A, Price G. Persistent psychotic symptoms following COVID-19 infection. *BJPsych Open.* 2020;6:e105. doi: 10.1192/bjo.2020.76.
7. Weerahandi H, Hochman KA, Simon E, Blaum C, Chodosh J, Duan E, et al. Post-discharge health status and symptoms in patients with severe COVID-19. *medRxiv.* 2020 Aug 14:2020.08.11.20172742. doi: 10.1101/2020.08.11.20172742.
8. Jany B. Pneumologische Ursachen für Thoraxschmerz [Pulmonary causes of chest pain]. *Internist (Berl).* 2017;58:22-28. doi: 10.1007/s00108-016-0169-9. [German].
9. Wächter C, Markus B, Schieffer B. Kardiologische Ursachen für Thoraxschmerz [Cardiac causes of chest pain]. *Internist (Berl).* 2017;58:8-21. doi: 10.1007/s00108-016-0165-0. [German].
10. Perrin R, Riste L, Hann M, Walther A, Mukherjee A, Heald A. Into the looking glass: Post-viral syndrome post COVID-19. *Med Hypotheses.* 2020;144:110055. doi: 10.1016/j.mehy.2020.110055.
11. Brewer G, Stratton K. Living with Chronic Fatigue Syndrome during lockdown and a global pandemic. *Fatigue: Biomedicine, Health & Behavior.* 2020;8:144-55. doi: 10.1080/21641846.2020.1827503.
12. Paolo G. Does COVID-19 cause permanent damage to olfactory and gustatory function? *Med Hypotheses.* 2020;143:110086. doi: 10.1016/j.mehy.2020.110086.
13. Anwar MM, Badawi AM, Eltablawy NA. Can the coronavirus infection penetrates the brain resulting in sudden anosmia followed by severe neurological disorders? *eNeurological Sci.* 2020;21:100290. doi: 10.1016/j.ensci.2020.100290.
14. Plantone D, Koudriavtseva T. Current and future use of chloroquine and hydroxychloroquine in infectious, immune, neoplastic, and neurological diseases: a mini-review. *Clin Drug Investig.* 2018;38:653-671. doi: 10.1007/s40261-018-0656-y.